



# ORLISSA

The first and only FDA-approved, hormone-free pill clinically proven to relieve moderate to severe endometriosis pain

Not an actual patient.

## USE

ORLISSA® (elagolix) is a prescription medicine used to treat moderate to severe pain associated with endometriosis. It is not known if ORLISSA is safe and effective in children.

## SAFETY CONSIDERATIONS

Do not use ORLISSA if you are pregnant, have osteoporosis or severe liver disease, take medicines called organic anion transporting polypeptide (OATP) 1B1 inhibitors that are known or expected to significantly increase the blood levels of elagolix (the active ingredient in ORLISSA), or have had a serious allergic reaction to ORLISSA or any of the ingredients in ORLISSA. ORLISSA does not prevent pregnancy. It may alter your period, so watch for other signs of pregnancy. Stop taking ORLISSA if you become pregnant. Ask your healthcare provider about proper birth control, as some may affect how ORLISSA works. ORLISSA may affect how some birth control works. ORLISSA can cause serious side effects, including bone loss, abnormal liver tests, suicidal thoughts or behaviors, and worsening mood. Talk to your healthcare provider right away if you notice changes such as jaundice, dark amber-colored urine, suicidal thoughts or actions, depression, or worsening mood.

**Please see Important Safety Information throughout and accompanying Consumer Brief Summary, or visit [www.rxabbvie.com/pdf/orilissa\\_pi.pdf](http://www.rxabbvie.com/pdf/orilissa_pi.pdf) for full Prescribing Information.**



**Orilissa**<sup>®</sup>  
elagolix tablets 150 mg  
200 mg



Not actual patients.

## DISCOVER ORILISSA

ORILISSA is not a painkiller, injection, or birth control. It's a hormone-free pill specifically proven to reduce moderate to severe endometriosis pain.

Doctors have already prescribed it to over 125,000 women.\*

We're so glad you want more info about ORILISSA. Take a look through this brochure, then ask your healthcare provider if ORILISSA could be right for you.

\*As of March 2023.

### IMPORTANT SAFETY INFORMATION

**What is the most important information I should know about ORILISSA?**

**ORILISSA may cause serious side effects, including:**

- **Bone Loss (decreased Bone Mineral Density [BMD])**  
While you are taking ORILISSA, your estrogen levels will be low. This can lead to BMD loss. If you have bone loss on ORILISSA, your BMD may improve after stopping ORILISSA, but may not recover completely. It is unknown if these bone changes could increase your risk for broken bones as you age. For this reason, your healthcare provider (HCP) may limit the length of time you take ORILISSA. Your HCP may order a DXA scan to check your BMD.
- **Effects on Pregnancy**  
**Do not take** ORILISSA if you are trying to become or are pregnant, as your risk for early pregnancy loss may increase. **If you think you are pregnant**, stop taking ORILISSA right away and call your HCP.

Please see Important Safety Information throughout and accompanying **Consumer Brief Summary**, or visit [www.rxabbvie.com/pdf/orilissa\\_pi.pdf](http://www.rxabbvie.com/pdf/orilissa_pi.pdf) for full Prescribing Information.

## PROVEN IN CLINICAL STUDIES

ORILISSA was studied in the largest endometriosis clinical trial program to date<sup>†</sup>:

1686  
WOMEN

AGES  
18 TO 49

WITH  
MODERATE  
TO SEVERE  
ENDO PAIN

ORILISSA was proven to relieve 3 common types of endo pain:

1. Painful periods
2. Pelvic pain in between periods
3. Pain with sex<sup>‡</sup>

<sup>†</sup>Evaluated in two 6-month clinical trials—952 women were treated with ORILISSA (475 with 150 mg once daily and 477 with 200 mg twice daily) and 734 were treated with placebo.

<sup>‡</sup>Only the ORILISSA 200 mg twice-daily dose was proven to work for pain with sex.

**When to expect feeling relief from endo pain:**

ORILISSA starts working in the body to lower estrogen after about 24 hours. However, pain relief will take longer. In clinical studies, ORILISSA was proven to provide relief at 3 months.

### IMPORTANT SAFETY INFORMATION (cont'd)

**What is the most important information I should know about ORILISSA? (cont'd)**

- **Effects on Pregnancy (cont'd)**  
ORILISSA may change your menstrual periods (irregular bleeding or spotting, a decrease in menstrual bleeding, or no bleeding at all), making it hard to know if you are pregnant. Watch for other signs of pregnancy, such as breast tenderness, weight gain, and nausea. ORILISSA does not prevent pregnancy. You will need to use effective hormone-free birth control (such as condoms or spermicide) while taking ORILISSA and for 28 days after stopping ORILISSA. Birth control pills that contain estrogen may make ORILISSA less effective. It is unknown how well ORILISSA works while on progestin-only birth control.

**Do not take ORILISSA if you:**

- Are pregnant, have osteoporosis, have severe liver disease, are taking medicines called organic anion transporting polypeptide (OATP) 1B1 inhibitors that are known or expected to significantly increase the blood levels of elagolix, the active ingredient in ORILISSA (ask your HCP if you are not sure if you are taking one of these medicines), or have had a serious allergic reaction to ORILISSA or any of the ingredients in ORILISSA. See the end of the Medication Guide for a complete list of ingredients in ORILISSA. Ask your HCP if you are not sure.

 **Orilissa**<sup>®</sup>  
elagolix tablets 150 mg  
200 mg

## HOW ORLISSA WORKS

ORLISSA is made for women with moderate to severe endometriosis pain.

It's a hormone-free pill that works by dialing down estrogen.



**ESTROGEN IS A HORMONE THAT'S NATURALLY PRODUCED BY YOUR BODY**




**ESTROGEN FUELS THE GROWTH OF PAINFUL ENDOMETRIOSIS LESIONS**



**ORLISSA COMES IN 2 HORMONE-FREE DOSES, EACH OF WHICH LOWERS ESTROGEN TO A DIFFERENT LEVEL**

Lower estrogen levels can help manage endo pain.

Watch the video to see ORLISSA in action



Visit [ORLISSA.com/How](https://www.Orilissa.com/How)

### IMPORTANT SAFETY INFORMATION (cont'd)

**What should I tell my HCP before taking ORLISSA?**

**Tell your HCP about all of your medical conditions, including if you:**

- Have or have had broken bones or other conditions that may cause bone problems; have or have had depression, mood problems, or suicidal thoughts or behavior; have liver problems; think you may be pregnant; or are breastfeeding or plan to be. It is unknown if ORLISSA passes into breast milk. Talk to your HCP about the best way to feed your baby if you take ORLISSA.

**Tell your HCP about all of the medicines you take,** including prescription and over-the-counter medicines, vitamins, and herbal supplements.

**Especially tell your HCP if you take** birth control that contains hormones. Your HCP may advise you to change your method of birth control.

**Please see Important Safety Information throughout and accompanying Consumer Brief Summary, or visit [www.rxabbvie.com/pdf/orilissa\\_pi.pdf](https://www.rxabbvie.com/pdf/orilissa_pi.pdf) for full Prescribing Information.**

## START MANAGING YOUR ENDOMETRIOSIS PAIN

ORLISSA is a daily, hormone-free pill available in 2 doses:

**ORLISSA 150 mg** One pill taken **once a day**, at about the same time each day, with or without food

**ORLISSA 200 mg** One pill taken **twice a day**, for a total of 400 mg a day, at about the same times each day, with or without food

Your healthcare provider will recommend the lowest effective dose based on your symptoms and treatment goals. Both the pain relief and any side effects you may experience could be different, depending on the dose you take and how your body responds. Because ORLISSA treats endometriosis pain by lowering estrogen, there are possible related side effects. Learn more about Important Safety Information for ORLISSA throughout this brochure.



It's important to **take ORLISSA every day**, exactly as your healthcare provider prescribed—**even if you're feeling better.**

Get more details on how to take ORLISSA, or what to do if you forget a dose, at [ORLISSA.com/about/how-to-take-ORLISSA](https://www.Orilissa.com/about/how-to-take-ORLISSA)

### IMPORTANT SAFETY INFORMATION (cont'd)

**What are the possible side effects of ORLISSA?**

**ORLISSA can cause serious side effects including:**

- **Suicidal thoughts, actions, or behavior, and worsening of mood. Call your HCP or get emergency medical help right away if you have any of these symptoms, especially if they are new, worse, or bother you:** thoughts about suicide or dying, attempts to commit suicide, new or worse depression or anxiety, or other unusual changes in behavior or mood. You or your caregiver should pay attention to any changes, especially sudden changes in your mood, behaviors, thoughts, or feelings.
- **Abnormal liver tests. Call your HCP right away if you have any of these signs and symptoms of liver problems:** yellowing of the skin or the whites of the eyes (jaundice), dark amber-colored urine, feeling tired, nausea and vomiting, generalized swelling, right upper stomach area pain, or bruising easily.



# SAVE ON ORILISSA

If you are a commercially insured patient, you may pay as little as \$5 a month or as little as \$15 per 3 month supply with the ORILISSA Savings Card.\*



Scan QR code for ORILISSA Savings Card or visit [ORILISSASavingsCard.com](https://ORILISSASavingsCard.com)



\*Eligibility: Available to patients with commercial insurance coverage for ORILISSA® (elagolix) who meet eligibility criteria. This co-pay assistance program is not available to patients receiving prescription reimbursement under any federal, state, or government-funded insurance programs (for example, Medicare [including Part D], Medicare Advantage, Medigap, Medicaid, TRICARE, Department of Defense, or Veteran Affairs programs) or where prohibited by law. Offer subject to change or termination without notice. Restrictions, including monthly maximums, may apply. This is not health insurance. **For full Terms and Conditions, visit [ORILISSASavingsCard.com](https://ORILISSASavingsCard.com) or call 1-800-ORILISSA for additional information.** To learn more about AbbVie's privacy practices and your privacy choices, visit <https://privacy.abbvie>

## IMPORTANT SAFETY INFORMATION (cont'd)

### What are the possible side effects of ORILISSA? (cont'd)

**The most common side effects of ORILISSA include:** hot flashes and night sweats, headache, nausea, difficulty sleeping, absence of periods, anxiety, joint pain, depression, and mood changes.

These are not all of the possible side effects of ORILISSA. This is the most important information to know about ORILISSA. For more information, talk to your HCP.

Take ORILISSA exactly as your HCP tells you. Tell your HCP if you have any side effect that bothers you or that does not go away. Call your doctor for medical advice about side effects.

**You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](https://www.fda.gov/medwatch) or call 1-800-FDA-1088.**

**If you are having difficulty paying for your medicine, AbbVie may be able to help. Visit [AbbVie.com/myAbbVieAssist](https://AbbVie.com/myAbbVieAssist) to learn more.**

**Please see Important Safety Information throughout and accompanying Consumer Brief Summary, or visit [www.rxabbvie.com/pdf/orilissa\\_pi.pdf](https://www.rxabbvie.com/pdf/orilissa_pi.pdf) for full Prescribing Information.**

abbvie

© 2024 AbbVie. All rights reserved.  
ORILISSA® and its design are  
registered trademarks of AbbVie Inc.  
US-ORIL-240004 March 2024

 **Orilissa®**  
elagolix tablets 150 mg  
200 mg